

# Study on Biological Characteristics of Ammonia-oxidizing Archaea in the Ocean

著者	李 沁潼
内容記述	この博士論文は内容の要約のみの公開（または一部非公開）になっています
year	2016
その他のタイトル	海洋性アンモニア酸化古細菌の生理生態特性に関する研究
学位授与大学	筑波大学 (University of Tsukuba)
学位授与年度	2015
報告番号	12102甲第7784号
URL	<a href="http://hdl.handle.net/2241/00144298">http://hdl.handle.net/2241/00144298</a>

**Study on Biological Characteristics of  
Ammonia-oxidizing Archaea in the Ocean**

January 2016

Qintong LI

**Study on Biological Characteristics of  
Ammonia-oxidizing Archaea in the Ocean**

A Dissertaion Submitted to  
the Graduate School of Life and Environmental Sciences,  
the University of Tsukuba  
in Partial Fulfillment of the Requirements  
for the Degree of Doctor of Philosophy in Biological Sciences  
(Doctoral Program in Bioindustrial Sciences)

Qintong LI

## Contents

Chapter 1 General introduction.....	1
1.1 Oceanic archaea .....	1
1.1.1 Taxonomy.....	1
1.1.2 Involvement of archaea in biogeochemical cycle.....	2
1.2 Global nitrogen cycle.....	3
1.2.1 Marine nitrogen cycle .....	3
1.2.2 Ammonia oxidation.....	4
1.3 Some open questions about ammonia-oxidizing archaea .....	4
1.3.1 Relative distribution of ammonia-oxidizing archaea and ammonia-oxidizing bacteria.....	4
1.3.2 Environmental factors affecting the distribution of ammonia-oxidizing archaea.....	5
1.3.3 Thaumarchaeota: autotrophic, heterotrophic or mixotrophic?.....	5
1.4 Oceanography .....	6
1.4.1 The Pacific sector of the Arctic Ocean.....	6
1.4.2 The North Pacific Ocean.....	7
1.4.3 The Suruga Bay, Japan.....	8
1.5 Objectives of this study.....	8
Chapter 2 Distribution and diversity of ammonia-oxidizing archaea in the Pacific sector of the Arctic Ocean .....	15
2.1 Introduction.....	15
2.2 Materials and methods .....	15
2.2.1 Sample collection.....	15
2.2.2 DNA extraction .....	16
2.2.3 Assay validation and optimization for real-time PCR (qPCR) .....	17
2.2.4 Real-time PCR (qPCR) analysis .....	18
2.2.5 Clone library construction, sequencing, and phylogenetic analysis of archaeal <i>amoA</i>	

gene .....	19
2.2.6 Statistical analysis .....	21
2.3 Results.....	21
2.3.1 Environment in the Pacific sector of the Arctic Ocean.....	21
2.3.2 Result of assay validation and optimization. ....	23
2.3.4 Vertical distributions of bacterial <i>amoA</i> gene .....	23
2.3.3 Vertical distributions of thaumarchaeal <i>amoA</i> gene in the Pacific sector of the Arctic Ocean .....	24
2.3.5 Vertical distributions of prokaryotic 16S rRNA genes .....	24
2.3.6 Vertical distributions of thaumarchaeal and bacterial <i>ureC</i> gene.....	25
2.3.7 Diversity of archaeal <i>amoA</i> gene at St. 154.....	26
2.4 Discussion .....	27
2.4.1 What determines the vertical distribution of AOA in the Pacific sector of the Arctic Ocean?.....	27
2.4.2 The relative distribution of AOA and AOB .....	30
2.4.3 AOA versus Thaumarchaeota .....	32
2.4.4 Possible ureolytic pathway for marine thaumarchaeota .....	34
2.4.5 About primer specificity .....	35
2.5 Conclusion .....	37
Chapter 3 Distribution and activity of ammonia-oxidizing archaea in the North Pacific Ocean .....	61
3.1 Introduction.....	61
3.2 Materials and method.....	61
3.2.1 Sample collection.....	61
3.2.2 Nucleic acid extraction .....	62
3.2.3 Reverse transcript PCR (RT-PCR) analysis .....	63
3.2.4 Real-time PCR (qPCR) analysis .....	64
3.3 Results.....	65

3.3.1 Environment in sampling sites in the North Pacific Ocean .....	65
3.3.2 Distribution and transcriptional activity of thaumarchaeal <i>amoA</i> in the North Pacific Ocean .....	66
3.3.3 Distribution and transcriptional activity of thaumarchaeal <i>ureC</i> in the North Pacific Ocean .....	67
3.3.4 Distribution of thaumarchaeal 16S rRNA gene in the North Pacific Ocean.....	68
3.4 Discussion .....	68
3.5 Conclusion .....	69
Chapter 4 Distribution and activity of ammonia-oxidizing archaea in the Suruga Bay, Japan	78
4.1 Introduction.....	78
4.2 Materials and method.....	78
4.2.1 Sample collection.....	78
4.2.3 Reverse transcript PCR (RT-PCR) analysis .....	79
4.2.4 Real-time PCR (qPCR) analysis .....	80
4.3 Results.....	81
4.3.1 Distribution and transcriptional activity of thaumarchaeal <i>amoA</i> in the Suruga Bay, Japan .....	81
4.3.2 Distribution and transcriptional activity of thaumarchaeal <i>ureC</i> in the North Pacific Ocean .....	81
4.4 Discussion .....	82
4.5 Conclusion .....	83
Chapter 5 Conclusions and perspectives.....	88
Acknowledgement .....	91
References .....	93

## Chapter 1 General introduction

### 1.1 Oceanic archaea

#### 1.1.1 Taxonomy

Since the concept of Archaea has been presented by Woese and Fox in 1977 (Woese and Fox, 1977), Archaea had been defined into two phyla, *Euryarchaeota* and *Crenarchaeota* (Woese *et al.*, 1990), and been considered as a kind of microorganism inhabiting in particular extreme environment. Among these two phyla, *Euryarchaeota* includes methanogenic archaea, highly halophilic archaea, thermophilic and acidophilic archaea as well as some hyperthermophilic archaea, while *Crenarchaeota* includes hyperthermophilic archaea. (Fig. 1-1)

Reported by DeLong in 1992, archaea was first proved to exist in a wide variety of environment, not only in some particular extreme environment (DeLong, 1992; Fuhrman *et al.*, 1992). Different from extremophilic archaea, marine mesophilic archaea can be divided into four phylogenetic groups: Marine Group I ~IV. Among these four groups, Marine Group I is a phylogenetic group belonging to *Crenarchaeota* (DeLong, 1992). And the other three groups (Marine Group II ~IV) are belonging to *Euryarchaeota* (DeLong, 1992; Fuhrman and Davis, 1997; López-García *et al.*, 2001). However, only few isolates in these groups are available yet, and their biological role in the environment remains to be fully elucidated.

Marine Group I is actually a distinct phylogenetic group which is phylogenetically close to the group of hyperthermophilic *Crenarchaeota* (DeLong, 1992). Recently, the concept of the third phylum of archaea: Thaumarchaeota (which comprises the phylogenetic group of Marine Group I) has been proposed (Brochier-Armanet *et al.*, 2008), and being supported by many researchers (Brochier-Armanet *et al.*, 2012; Pester *et al.*, 2011; Spang *et al.*, 2010; Walker *et al.*, 2010). Organisms in this phylum are generally mesophilic, and globally distributed and are found in high numbers in marine and freshwater environments, soils, and sediments.

Marine Group I and Marine Group II are the most abundant groups existing in the global ocean among marine archaea (DeLong *et al.*, 1999; Karner *et al.*, 2001). And Marine Group I is one of the main members of planktonic prokaryotes, accounting for 40~50% of

bathypelagic prokaryotes, and 20% of total oceanic prokaryotes (Karner *et al.*, 2001). Marine Group II are dominant in euphotic zone in temperate oceans, and existing in a variety of environment including the Antarctic Ocean (Massana *et al.*, 1998, 2000; Murrey *et al.*, 1999).

#### 1.1.2 Involvement of archaea in biogeochemical cycle

The biogeochemistry of our planet is shaped by the microbes-driven biological fluxes of the elements, particularly the six major building blocks of life: hydrogen, carbon, nitrogen, sulfur, oxygen and phosphorus. Since the concept 'biogeochemistry' has been proposed from 1926, only bacteria and eukarya were considered contribute to these nutrients cycles. The domain archaea, which had been thought to be habit only in extreme environments, were first described in the biogeochemical cycle until 50 years later (Woese and Fox, 1977). The role and significance of archaea started to be realized since early 1990, until the ubiquity and diversity of oceanic planktonic archaea has been recognized (DeLong *et al.*, 1992; Fuhrman *et al.*, 1992). Using molecular biological, biochemical and isotope-based methods, evidences began to accumulate proving that archaea are important contributors in carbon and nitrogen cycling.

#### **Archaea in the nitrogen cycle**

In the nitrogen cycle, archaea get involved in the fixation of dinitrogen gas, as the methanogenic archaea has the ability to fix dinitrogen gas; archaea also participate in denitrification process, as a few cultivated archaea are capable of denitrification. However, the most important role of archaea in the nitrogen cycle is they can aerobically oxidize ammonia into nitrite, and thus recently archaea has been recognized as the main driver of nitrification in various ecosystems (i.e. reviewed by Offre *et al.*, 2013).

#### **Archaea in the carbon cycle**

Many cultivated representatives of Crenarchaeota, Thaumarchaeota and Euryarchaeota are capable of autotrophic growth, as they assimilate carbon from oxidized inorganic compounds such as carbon dioxide or bicarbonate (Berg *et al.*, 2010). Nevertheless, many autotrophic members of the Archaea can grow mixotrophically, as they can coassimilate small organic compounds under suitable conditions or switch between an autotrophic and a heterotrophic



lifestyle (Kletzin, 2007).

Methane, a major greenhouse gas but also an important source of energy for humans, is the predominant hydrocarbon in earth's atmosphere. Methanogenic archaea, affiliated with Euryarchaeota, are strict anaerobes that produce methane as the major product of their energy conserving metabolism. Archaea dominate the biogenic production of methane, but also key to the oxidation of this important hydrocarbon. Some close relatives of methanogenic archaea play a critical role in controlling the emissions of methane, by oxidizing methane back to CO<sub>2</sub>. The methane-oxidizing archaea are also strict anaerobes and affiliated to Euryarchaeota. (i.e. reviewed by Offre et al., 2013)

## 1.2 Global nitrogen cycle

### 1.2.1 Marine nitrogen cycle

Nitrogen is biologically limiting element and an essential nutrient element to all living organisms in the planet. Although ca. 79% of the air is consist of nitrogen gas (N<sub>2</sub>), this stable form of nitrogen cannot be utilized by most living organisms because the triple bound structure is difficult and energetically cost to break. Thus, the fixed form of nitrogen, referring to any reactive form of nitrogen (such as nitrate, nitrite and ammonia...), are required by most living organisms (such as algae, and microbes...). The processes governing transitions from one form of nitrogen to another are all part of the nitrogen cycle. (Fig. 1-2)

At global scale, biological nitrogen fixation is the largest source of nitrogen to the ocean, while the anaerobic microbial processes are the main export of nitrogen losses (Francis et al., 2005; Galloway, et al., 2004; Gruber et al., 1997). Nevertheless, the biological nitrogen fixation and anaerobic nitrogen losses are ultimately connected by nitrification, the microbe-driven oxidation from ammonium to nitrite and nitrate. Every year, ca.  $4 \times 10^{11}$  kg of nitrogen cycles through the ocean each year (Francis et al., 2005; Galloway, et al., 2004), and nearly all of this nitrogen has to be nitrified at least once (Karl, 2002). Given that nitrification controls the global nitrogen balance, this process is one of the most important ocean processes in global marine nitrification.

### 1.2.2 Ammonia oxidation

Ammonia oxidation is the first step and rate-limiting step of nitrification, the microbially mediated pathway for the conversion of ammonia to nitrite. It has long been considered that ammonia oxidation is conducted only by microorganisms from the domain *Bacteria*: the betaproteobacteria ( $\beta$ - Ammonia-oxidizing Bacteria;  $\beta$ -AOB) from the genus *Nitrosomonas* and *Nitrospira* (Purkhold et al., 2000; 2003; Beaumont et al., 2004; Taylor and Bottomley, 2006) and the gammaproteobacteria ( $\gamma$ - Ammonia-oxidizing Bacteria;  $\gamma$ -AOB) from the genus *Nitrosococcus* (Ward and O'Mullan, 2002). However, these two groups of AOB often comprise only 0.1% of the bacterial assemblages (Ward et al., 2000). Until 2004, based on the discovery of a unique ammonia monooxygenase gene (*amoA* gene) on archaea domain (Vent et al., 2004), the involvement of archaea in nitrification was first suggested. Since then, the ubiquity and diversity of ammonia-oxidizing archaea (AOA) has been increasingly recognized, and these ammonia-oxidizing archaea were found specifically associated with the pelagic non-thermophilic marine thaumarchaeota (Beman et al., 2006; Agogue et al., 2008; Cafferey et al., 2007; Coolen et al., 2007; Lam et al., 2007; Santoro et al., 2008). In contrast to the low abundance of AOB reported in various eco-systems, thaumarchaeota are ubiquitous and abundant in the ocean (DeLong et al., 1994; Stein and Simon, 1996; Karner et al., 2001; Herndl et al., 2005; Ingalls et al., 2006), and often account for ca. 20% of all prokaryotic cells in the global ocean (Karner et al., 2001). Thus, the archaea-driven nitrification may play an important role in global oceanic nitrogen cycle, whereas the potential relative contribution of bacterial nitrification and archaeal nitrification in natural waters still remain to be fully elucidated.

### 1.3 Some open questions about ammonia-oxidizing archaea

#### 1.3.1 Relative distribution of ammonia-oxidizing archaea and ammonia-oxidizing bacteria

The wide distribution of ammonia-oxidizing archaea (AOA), affiliated with Thaumarchaeota, across a variety of aquatic environments is now well established through reports based on abundance of the gene-encoding archaeal ammoniamonooxygenase  $\alpha$ -subunit (*amoA*) in

oceanic waters (Beman et al., 2006; Agogue et al., 2008; Caffrey et al., 2007; Coolen et al., 2007; Lam et al., 2007; Santoro et al., 2008). Nevertheless, the relative contribution of AOA versus AOB remains unclear and factors that regulate ammonia oxidizers' activity and diversity in aquatic ecosystems have not yet been fully elucidated. Geochemical-based approaches and molecular-based approaches targeting both cultured strains and uncultured environmental samples indicate that AOA are very different from AOB, their ammonia-oxidizing bacterial counterparts. Physiological differences between AOA and AOB have been pointed to aspects such as ammonia affinity (Martens-Habben et al., 2009), substrate range (Könneke et al., 2005; Hallam et al., 2006; Ouverney et al., 2000), and metabolic pathway (Hallam et al., 2006b; Berg et al., 2011; Blainey et al., 2011; Park et al., 2010; Arp et al., 2007). These physiological differences are expected to be vital factors and be reflected in their niche segregation and the observed dynamics of their relative distributions.

### 1.3.2 Environmental factors affecting the distribution of ammonia-oxidizing archaea

Pelagic marine Thaumarchaeota are typically most abundant below ~100 m depth in the water column (DeLong, 1992; Fuhrman et al., 1992; Massana et al., 1997; Karner et al., 2001; Mincer et al., 2007; Church et al., 2010; Santoro et al., 2010). Previous studies have pointed to environmental factors such as salinity, light, temperature, ammonium, oxygen, and sulfide as major determinants of the distribution of ammonia oxidizing archaea (e.g., Murray et al., 1999; Caffrey et al., 2007; Santoro et al., 2008; Bernhard et al., 2010; Gubry-Rangin et al., 2010), however, the results are often contradictory. For most previous studies, the distributions of AOA have been investigated on the euphotic and mesopelagic zone, where ammonia concentrations are much higher than in waters below 1000 m depth. Thus, a thorough study targeting throughout water column on a large-scale sampling are of great importance for better understand the major environmental regulators of the distribution of AOA.

### 1.3.3 Thaumarchaeota: autotrophic, heterotrophic or mixotrophic?

The natural radiocarbon content archaeal lipids indicated that thaumarchaeota in ocean

waters predominately grow autotrophically (Ingalls et al., 2006; Pearson et al., 2001; Wutcher et al., 2003). Modeled estimates of archaeal biomass production in deep waters range from 0.7 to 0.8 Gt C/year, suggesting thaumarchaeota might contribute ca. 1% of the annual primary production in the ocean (Herndl et al., 2005; Ingalls et al., 2006). However, a potential for mixotrophic growth has been reported from analyses of the genome of *Nitrosopumilus maritimus* SCM1 and *Ca. Cenarchaeum symbiosum* (Hallam et al., 2006a; 2006b; Walker et al., 2010). And a recent research revealed that two new marine AOA isolates *Nitrosopumilus maritimus* PS0 and HAC1 were obligate mixotrophs that rely on uptake and assimilation of organic carbon compounds (Qin et al., 2014).

Additionally, a urea-utilization mechanism hypothesized to explain thaumarchaeal nitrification, which has been recently suggested by urea uptake experiments using seawater samples (Alonso-Sáez et al., 2012) and isolated marine AOA strains (Qin et al., 2014), shows the presumed significance of an urelytic mechanism under low-energy conditions. However, other studies have shown that transcripts of the functional gene in the urea-utilization mechanism, the *ureC* gene, are not common in surface waters of the Canadian Arctic (Pedneault, et al., 2014) and Marennes-Oléron estuary (Hugoni et al., 2015). Therefore, we still lack sufficient evidence to show the possibility and significance of this urelytic mechanism.

## 1.4 Oceanography

### 1.4.1 The Pacific sector of the Arctic Ocean

The Arctic Ocean (Fig. 1-3), located in the northernmost part of the world, is one of the coldest areas in the world. However, the Arctic systems can be rich and diverse habitats for marine life in spite of the extreme cold environment (Grebmeier et al. 2006, 2010). Biological cycling processes and biological communities are directly influenced by changing sea ice extent, light availability, seawater hydrography (such as nutrients, salinity, temperature, density), currents, and water column production (Grebmeier and Maslowski et al., 2014). The retreating sea ice and warming temperature have dramatic impacts on the biological system, such as changes in overlying primary production, altering organic carbon transformation,

pelagic-benthic production and community structure that could have further effects to higher trophic levels (Grebmeier and Maslowski et al., 2014).

The Pacific sector of the Arctic Ocean is experiencing the greatest seasonal retreat and thinning of sea ice in the Arctic (Grebmeier and Maslowski et al., 2014). In the Pacific sector of the Arctic Ocean, Pacific water transiting across the wide Bering, Chukchi and the eastern portion of the Beaufort Sea, is a major driving force for the physical environmental state, ice extent and thickness, productivity and carbon transport in the Amerasian Arctic (Grebmeier and Maslowski et al., 2014). In this region, the Bering Strait is a major 'gate' and a pathway for fluxes of biological organisms and organic carbon (Grebmeier et al. 2006; Grebmeier, 2012 ; Walsh et al., 2004). Recent estimates on the Bering Strait indicated that freshwater in Bering Strait may provide ca. 40% of the total freshwater input to the Arctic Ocean (Woodgate and Aagaard, 2005). The nutrient rich Pacific waters transit through the Bering Strait, and influence the thermohaline circulation in the Arctic Ocean. Winter and summer Pacific Water types play different roles in the transport of heat, fresh water, nutrients, carbon and biological organisms through the Bering Strait. The warm Pacific summer water is a source of heat and particularly impact the Chukchi Sea area; while the cold Pacific winter water influence the halocline formation (Shimada et al., 2006).

#### 1.4.2 The North Pacific Ocean

The Pacific Ocean (Fig. 1-4) is the largest oceanic division on earth, which connects the Arctic Ocean in the north and the Antarctic Ocean in the south. The Pacific Ocean also extends from Asia and Australia in the west and the Americas in the east, covering 165.25 million square kilometers in area and representing ca. 50% of the global earth's oceanic water (<http://global.britannica.com/place/Pacific-Ocean>), and thus has an important role in controlling the global climate. The equator divides the Pacific Ocean into North Pacific Ocean and South Pacific Ocean. The motion of Pacific waters is generally clockwise in the North Pacific Gyre and counter-clockwise in the South Pacific Gyre. The North Equatorial Current, driven westward along latitude 15°N by the trade winds, turns north near the Philippines to

become the warm Kuroshio Current.

In the North Pacific Ocean, most of it is covered by the North Pacific Gyre, which locates between the equator and 50°N latitude. The northern part of North Pacific Gyre is the North Pacific Current, which is a slow warm water current that flows west to east between 40°N and 50°N. When it flows near Americas, this current splits into the southward California Current and northward Alaska Current. The southward California Current then moves to the west, joining the North Equatorial Current and circling in the North Pacific Gyre, while the northward Alaska Current then joining a smaller gyre in North Pacific Ocean, the Subpolar Gyre. The counterclockwise Subpolar Gyre forms at high latitudes and circulate along the Siberian Pacific coast and the subpolar areas in the North Pacific Ocean.

#### 1.4.3 The Suruga Bay, Japan

The Suruga Bay (Fig. 1-5) located on the Pacific coast of Honshu in Shizuoka Prefecture, Japan. As the nearby Mount Fuji (Japan's loftiest peak, 3776 m) rises from the 2500 m depth Suruga Trough running up the middle of the bay, Suruga Bay is the deepest nearshore bay in Japan. The Suruga Bay receives numerous fresh water inflows from rivers such as the Oi River, Fuji River, and Abe river from the western side; and the Kano River from the eastern side of Suruga Bay (<http://www.isewan-db.go.jp/index.asp>). Although the Suruga Bay is open to the Pacific Ocean to the south, it is protected from oceanic waves by Izu Peninsula. Thus, Suruga Bay has been one of the most important deep-sea fishery spot in Japan, famous for its rich fishery species diversity. For reasons above, the Suruga Bay is considered as a perfect research spot for studies on deep-sea organisms in temperate zone.

#### 1.5 Objectives of this study

In order to describe the distribution characteristics of ammonia oxidizing archaea, associated with dynamics of community structure and environmental variables, the abundance, diversity and activity of AOA were compared in three different ocean areas: the Pacific sector of the Arctic Ocean, the North Pacific Ocean, as well as the Suruga Bay in Japan. A total of 173

samples throughout water columns across these three oceans were analyzed based on molecular technologies targeting functional genes and 16S rRNA genes. This study presents a comprehensive description on the distribution characteristics of AOA, which is fundamental to a better understanding on the physiology of AOA, and provide valuable basis for perspectives of global microbial nitrification.




Fig.1-1 Evolutionary relationships of Archaea, Bacteria, and Eucarya. (modified from DeLong et al., 1998)






Fig.1-2 Marine nitrogen cycle (modified from <http://bit.ly/wC3xZo>)




Fig. 1-3 Map and bathymetry of the Arctic Ocean. (Edited based on The International Bathymetric Chart of the Arctic Ocean (IBCAO), Jakobsson et al., 2012)




Fig. 1-4 Map and oceanography of the Pacific Ocean (Modified based on graph from <http://www.seos-project.eu/modules/oceancurrents/oceancurrents-c02-p03.html>)




Fig. 1-5 Map and major rivers inflow of the Suruga Bay, Japan (Modified based on graph from <http://www.isewan-db.go.jp/suruga-gaiyo/A2c.asp>)

## **Acknowledgement**

I would like to special thank Prof. Motoo Utsumi for these five years' scientific guidance and instructions, as well as his generous help and support in my studies and life in Japan. Every progress and success I ever achieved during these five years cannot be separated with Prof. Utsumi's help and guidance.

I sincerely thank Prof. Kouji Nakamura, Prof. Toshiaki Nakajima and Prof. Yingnan Yang for reviewing this work and giving me precious suggestion, from which I benefit a lot to improve this work. I also really appreciate the valuable guidance and instructions from Prof. Norio Sugiura, Prof. Yang Yingnan, and Prof. Zhang Zhenya throughout my Ph.D. course.

Dr. Chie Amano-Sato and Mr. Shohei Akiyama contribute to this work in the Pacific Arctic sample collection. Dr. Amano-Sato and Mr. Akiyama also taught me a lot as senior lab members, their inspiring comments and kindly assistance are of great importance to this work and to myself. I'm grateful for the help of Mr. Seiya Takahashi, Mr. Toshiaki Eto, Dr. Phan Thann Luu, Ms. Ayako Kanazawa, Ms. Azusa Hashimoto and Mr. Shunichi Ohyama in sample collection in Suruga Bay. I would also remember the generous help of Dr. Minoru Ijichi from AORI, University of Tokyo, for teaching me real-time PCR technology and being supportive in my research plan.

I deeply thank captain and crew from RV 'Mirai' for their help in collecting samples in the Pacific sector of the Arctic Ocean as well as North Pacific, Captain Hiroshi Hasegawa and his son from RV 'Choganemaru' for their help in collecting samples in Suruga Bay, and the technicians from Global Ocean Development and Marine Works

Japan for sample collection and analysis.

I also greatly thank my laboratory members and friends, Dr. Gao Yu, Dr. Wang Xin, Mr. Cui Hailong, and all the other lab members for being together with me, sharing happiness and being supportive in every detail of my studying life. My lab members and friends left me a really joyful and memorable five-year.

Last but not least, I would like to thank my lovely family: my father Li Wei, my mother Dr. Wang Jin, my husband Du Ye, my cousin Hu Junzi, as well as my grandparents, my uncles and my aunties. My whole family fully motivate, support and encourage me to be a better scientist, and I will move forward in this career with their generous support.



## References

- Agogué H., Brink, M., Dinasquet, J., & Herndl, G. J. (2008). Major gradients in putatively nitrifying and non-nitrifying Archaea in the deep North Atlantic. *Nature*, 456 (7223), 788-791.
- Alonso-Sáez, L., Waller, A. S., Mende, D. R., Bakker, K., Farnelid, H., Yager, P. L., Lovejoy, C., Tremblay, J., Potvin, M., Heinrich, F., Estrada, M., Riemann, L., Bork, P., Pedró-Alí C., & Bertilsson, S. (2012). Role for urea in nitrification by polar marine Archaea. *Proceedings of the National Academy of Sciences*, 109 (44), 17989-17994.
- Amano-Sato, C., Akiyama, S., Uchida, M., Shimada, K., & Utsumi, M. (2013). Archaeal distribution and abundance in water masses of the Arctic Ocean, Pacific sector. *Aquatic microbial ecology*, 69 (2), 101-112.
- Arp, D. J., Chain, P. S., & Klotz, M. G. (2007). The impact of genome analyses on our understanding of ammonia-oxidizing bacteria\*. *Annu. Rev. Microbiol.*, 61, 503-528.
- Arrigo, K. R., van Dijken, G., & Pabi, S. (2008). Impact of a shrinking Arctic ice cover on marine primary production. *Geophysical Research Letters*, 35 (19).

Arrigo, K. R., Perovich, D. K., Pickart, R. S., Brown, Z. W., van Dijken, G. L., Lowry, K. E., ... & Swift, J. H. (2012). Massive phytoplankton blooms under Arctic sea ice. *Science*, 336 (6087), 1408-1408.

Beaumont, H. J., Lens, S. I., Reijnders, W. N., Westerhoff, H. V., & Van Spanning, R. J. (2004). Expression of nitrite reductase in *Nitrosomonas europaea* involves NsrR, a novel nitrite - sensitive transcription repressor. *Molecular microbiology*, 54 (1), 148-158.

Beman, J. M., & Francis, C. A. (2006). Diversity of ammonia-oxidizing archaea and bacteria in the sediments of a hypernutrified subtropical estuary: Bahia del Tobari, Mexico. *Applied and Environmental Microbiology*, 72 (12), 7767-7777.

Beman, J. M., Popp, B. N., & Francis, C. A. (2008). Molecular and biogeochemical evidence for ammonia oxidation by marine Crenarchaeota in the Gulf of California. *The ISME Journal*, 2 (4), 429-441.

Beman, J. M., Popp, B. N., & Alford, S. E. (2012). Quantification of ammonia oxidation rates and ammonia-oxidizing archaea and bacteria at high resolution in the Gulf of California and eastern tropical North Pacific Ocean. *Limnology and Oceanography*, 57 (3), 711-726.



Berg, I. A., Kockelkorn, D., Ramos-Vera, W. H., Say, R. F., Zarzycki, J., Hügler, M., Alber, B. E., & Fuchs, G. (2010). Autotrophic carbon fixation in archaea. *Nature Reviews Microbiology*, 8 (6), 447-460.

Berg, I. A. (2011). Ecological aspects of the distribution of different autotrophic CO<sub>2</sub> fixation pathways. *Applied and environmental microbiology*, 77 (6), 1925-1936.

Bernhard, A. E., Landry, Z. C., Blevins, A., José, R., Giblin, A. E., & Stahl, D. A. (2010). Abundance of ammonia-oxidizing archaea and bacteria along an estuarine salinity gradient in relation to potential nitrification rates. *Applied and Environmental Microbiology*, 76 (4), 1285-1289.

Blainey, P. C., Mosier, A. C., Potanina, A., Francis, C. A., & Quake, S. R. (2011). Genome of a low-salinity ammonia-oxidizing archaeon determined by single-cell and metagenomic analysis. *PLoS One*, 6 (2), e16626.

Bouskill, N. J., Eveillard, D., Chien, D., Jayakumar, A., & Ward, B. B. (2012). Environmental factors determining ammonia - oxidizing organism distribution and diversity in marine environments. *Environmental microbiology*, 14 (3), 714-729.

Braak, C. J. F., & Šmilauer, P. (2012). *CANOCO reference manual and user's guide: software for ordination (version 5.0)*. Biometris.

Brochier-Armanet, C., Boussau, B., Gribaldo, S., & Forterre, P. (2008). Mesophilic Crenarchaeota: proposal for a third archaeal phylum, the Thaumarchaeota. *Nature Reviews Microbiology*, 6 (3), 245-252.

Brochier-Armanet, C., Gribaldo, S., & Forterre, P. (2012). Spotlight on the Thaumarchaeota. *ISME Journal-International Society for Microbial Ecology*, 6 (2), 227.

Caffrey, J. M., Bano, N., Kalanetra, K., & Hollibaugh, J. T. (2007). Ammonia oxidation and ammonia-oxidizing bacteria and archaea from estuaries with differing histories of hypoxia. *The ISME journal*, 1 (7), 660-662.

Cai, W. J., Bates, N. R., Guo, L., Anderson, L. G., Mathis, J. T., Wanninkhof, R., ... & Semiletov, I. P. (2014). Carbon Fluxes Across Boundaries in the Pacific Arctic Region in a Changing Environment. In *The Pacific Arctic Region* (pp. 199-222). Springer Netherlands.

Christman, G. D., Cottrell, M. T., Popp, B. N., Gier, E., & Kirchman, D. L. (2011). Abundance, diversity, and activity of ammonia-oxidizing prokaryotes in the coastal Arctic Ocean in summer and winter. *Applied and environmental microbiology*, 77 (6), 2026-2034.

Church, M. J., Wai, B., Karl, D. M., & DeLong, E. F. (2010). Abundances of crenarchaeal *amoA* genes and transcripts in the Pacific Ocean. *Environmental Microbiology*, 12 (3), 679-688.

Coolen, M. J., Abbas, B., Van Bleijswijk, J., Hopmans, E. C., Kuypers, M. M., Wakeham, S. G., & Sinninghe Damste, J. S. (2007). Putative ammonia - oxidizing Crenarchaeota in suboxic waters of the Black Sea: a basin - wide ecological study using 16S ribosomal and functional genes and membrane lipids. *Environmental Microbiology*, 9 (4), 1001-1016.

De Corte, D., Yokokawa, T., Varela, M. M., Agogue H., & Herndl, G. J. (2009). Spatial distribution of Bacteria and Archaea and *amoA* gene copy numbers throughout the water column of the Eastern Mediterranean Sea. *The ISME journal*, 3 (2), 147-158.

De la Torre, J. R., Walker, C. B., Ingalls, A. E., Könneke, M., & Stahl, D. A. (2008). Cultivation of a thermophilic ammonia oxidizing archaeon synthesizing crenarchaeol. *Environmental Microbiology*, 10 (3), 810-818.

DeLong, E. F. (1992). Archaea in coastal marine environments. *Proceedings of the National Academy of Sciences*, 89 (12), 5685-5689.

DeLong, E. F., Wu, K. Y., Prézelin, B. B., & Jovine, R. V. (1994). High abundance of Archaea in Antarctic marine picoplankton. *Nature*, 371 (6499), 695-697.

DeLong, E. F., King, L. L., Massana, R., Cittone, H., Murray, A., Schleper, C., & Wakeham, S. G. (1998). Dibiphytanyl ether lipids in nonthermophilic crenarchaeotes. *Applied and Environmental Microbiology*, 64 (3), 1133-1138.

DeLong, E. F., Taylor, L. T., Marsh, T. L., & Preston, C. M. (1999). Visualization and enumeration of marine planktonic archaea and bacteria by using polyribonucleotide probes and fluorescent in situ hybridization. *Applied and Environmental Microbiology*, 65 (12), 5554-5563.

Francis, C. A., Roberts, K. J., Beman, J. M., Santoro, A. E., & Oakley, B. B. (2005). Ubiquity and diversity of ammonia-oxidizing archaea in water columns and sediments of the ocean. *Proceedings of the National Academy of Sciences of the United States of America*, 102 (41), 14683-14688.

Francis, C. A., Beman, J. M., & Kuypers, M. M. (2007). New processes and players in the nitrogen cycle: the microbial ecology of anaerobic and archaeal ammonia oxidation. *The ISME journal*, 1 (1), 19-27.

French, E., Kozlowski, J. A., Mukherjee, M., Bullerjahn, G., & Bollmann, A. (2012).

Ecophysiological characterization of ammonia-oxidizing archaea and bacteria from freshwater. *Applied and environmental microbiology*, 78 (16), 5773-5780.

Fuhrman, J. A. (1992). Novel major archaeobacterial group from marine plankton. *Nature*, 356 (6365), 148-149.

Fuhrman, J. A., & Davis, A. A. (1997). Widespread Archaea and novel Bacteria from the deep sea as shown by 16S rRNA gene sequences. *Oceanographic Literature Review*, 9 (44), 1025.

Galand, P. E., Lovejoy, C., Hamilton, A. K., Ingram, R. G., Pedneault, E., & Carmack, E. C. (2009). Archaeal diversity and a gene for ammonia oxidation are coupled to oceanic circulation. *Environmental microbiology*, 11 (4), 971-980.

Galloway, J. N., Dentener, F. J., Capone, D. G., Boyer, E. W., Howarth, R. W., Seitzinger, S. P., Asner, G. P., Cleveland, C. C., Green, P. A., Holland, E. A., Karl, D. M., Michaels, A. F., Porter, J. H., Townsend, A. R., & Vöosmarty, C. J. (2004). Nitrogen cycles: past, present, and future. *Biogeochemistry*, 70 (2), 153-226.

Grebmeier, J. M., Cooper, L. W., Feder, H. M., & Sirenko, B. I. (2006). Ecosystem dynamics of the Pacific-influenced northern Bering and Chukchi Seas in the Amerasian Arctic. *Progress in Oceanography*, 71 (2), 331-361.

Grebmeier, J. M., Moore, S. E., Overland, J. E., Frey, K. E., & Gradinger, R. (2010). Biological response to recent Pacific Arctic sea ice retreats. *Eos, Transactions American Geophysical Union*, 91 (18), 161-162.

Grebmeier, J. M. (2012). Shifting patterns of life in the Pacific Arctic and Sub-Arctic seas. *Marine Science*, 4, DOI: 10.1146/annurev-marine-120710-100926

Grebmeier, J. M., & Maslowski, W. (2014). The Pacific Arctic Region: An Introduction. In *The Pacific Arctic Region* (pp. 1-15). Springer Netherlands.

Gruber, N., & Sarmiento, J. L. (1997). Global patterns of marine nitrogen fixation and denitrification. *Global Biogeochemical Cycles*, 11 (2), 235-266.

Gubry-Rangin, C., Nicol, G. W., & Prosser, J. I. (2010). Archaea rather than bacteria control nitrification in two agricultural acidic soils. *FEMS Microbiology Ecology*, 74 (3), 566-574.

Hallam, S. J., Konstantinidis, K. T., Putnam, N., Schleper, C., Watanabe, Y. I., Sugahara, J., Preston, C., de la Torre, J., Richardson, P. M., & DeLong, E. F. (2006). Genomic analysis of the uncultivated marine crenarchaeote *Cenarchaeum symbiosum*. *Proceedings of the National Academy of Sciences*, 103 (48), 18296-18301.

Hallam, S. J., Mincer, T. J., Schleper, C., Preston, C. M., Roberts, K., Richardson, P. M., & DeLong, E. F. (2006). Pathways of carbon assimilation and ammonia oxidation suggested by environmental genomic analyses of marine Crenarchaeota.

Harrison, W. G., Head, E. J. H., Conover, R. J., Longhurst, A. R., & Sameoto, D. D. (1985). The distribution and metabolism of urea in the eastern Canadian Arctic. *Deep Sea Research Part A. Oceanographic Research Papers*, 32 (1), 23-42.

Herfort, L., Schouten, S., Abbas, B., Veldhuis, M. J., Coolen, M. J., Wuchter, C., Boon, J. P., Herndl, G. J., & Damsté J. S. S. (2007). Variations in spatial and temporal distribution of Archaea in the North Sea in relation to environmental variables. *FEMS microbiology ecology*, 62 (3), 242-257.

Herndl, G. J., Reinthaler, T., Teira, E., van Aken, H., Veth, C., Pernthaler, A., & Pernthaler, J. (2005). Contribution of Archaea to total prokaryotic production in the deep Atlantic Ocean. *Applied and Environmental Microbiology*, 71 (5), 2303-2309.

Hornek, R., Pommerening-Röser, A., Koops, H. P., Farnleitner, A. H., Kreuzinger, N., Kirschner, A., & Mach, R. L. (2006). Primers containing universal bases reduce multiple amoA gene specific DGGE band patterns when analysing the diversity of beta-ammonia oxidizers in the environment. *Journal of microbiological methods*, 66 (1), 147-155.

Hugoni, M., Agogué H., Taib, N., Domaizon, I., Moné A., Galand, P. E., ... & Mary, I. (2015). Temporal Dynamics of Active Prokaryotic Nitrifiers and Archaeal Communities from River to Sea. *Microbial ecology*, 1-11.

Ingalls, A. E., Shah, S. R., Hansman, R. L., Aluwihare, L. I., Santos, G. M., Druffel, E. R., & Pearson, A. (2006). Quantifying archaeal community autotrophy in the mesopelagic ocean using natural radiocarbon. *Proceedings of the National Academy of Sciences*, 103 (17), 6442-6447.

Jakobsson, M., Mayer, L., Coakley, B., Dowdeswell, J. A., Forbes, S., Fridman, B., Hodnesdal, H., Noormets, R., Pedersen, R., Rebescó, M., Schenke, H. W., Zarayskaya, Y., Accettella, D., Armstrong, A., Anderson, R. M., Bienhoff, P., Camerlenghi, A., Church, I., Edwards, M., Gardner, J. V., Hall, J. K., Hell, B., Hestvik, O., Kristoffersen, Y., Marcussen, C., Mohammad, R., Mosher, D., Nghiem, S. V., Pedrosa, M. T., Travaglini, P. G., & Weatherall, P. (2012). The international



bathymetric chart of the Arctic Ocean (IBCAO) version 3.0. *Geophysical Research Letters*, 39 (12).

Jung, M. Y., Park, S. J., Min, D., Kim, J. S., Rijpstra, W. I. C., Damsté J. S. S., Kim, G. J., Madsen, E. L., & Rhee, S. K. (2011). Enrichment and characterization of an autotrophic ammonia-oxidizing archaeon of mesophilic crenarchaeal group I. 1a from an agricultural soil. *Applied and environmental microbiology*, 77 (24), 8635-8647.

Kalanetra, K. M., Bano, N., & Hollibaugh, J. T. (2009). Ammonia - oxidizing Archaea in the Arctic Ocean and Antarctic coastal waters. *Environmental microbiology*, 11 (9), 2434-2445.

Karl, D., Michaels, A., Bergman, B., Capone, D., Carpenter, E., Letelier, R., Lipschultz, F., Paerl, H., Sigman, D., & Stal, L. (2002). *Dinitrogen fixation in the world's oceans* (pp. 47-98). Springer Netherlands.

Karner, M. B., DeLong, E. F., & Karl, D. M. (2001). Archaeal dominance in the mesopelagic zone of the Pacific Ocean. *Nature*, 409 (6819), 507-510.

Kletzin, A. (2007). Metabolism of inorganic sulfur compounds in Archaea. In *Archaea. Evolution, Physiology, and Molecular Biology* (pp.261-274). John Wiley & Sons.

Könneke, M., Bernhard, A. E., Jos é R., Walker, C. B., Waterbury, J. B., & Stahl, D. A. (2005). Isolation of an autotrophic ammonia-oxidizing marine archaeon. *Nature*, 437 (7058), 543-546.

Konstantinidis, K. T., Braff, J., Karl, D. M., & DeLong, E. F. (2009). Comparative metagenomic analysis of a microbial community residing at a depth of 4,000 meters at station ALOHA in the North Pacific subtropical gyre. *Applied and environmental microbiology*, 75 (16), 5345-5355.

Labrenz, M., Sintes, E., Toetzke, F., Zumsteg, A., Herndl, G. J., Seidler, M., & Jürgens, K. (2010). Relevance of a crenarchaeotal subcluster related to Candidatus Nitrosopumilus maritimus to ammonia oxidation in the suboxic zone of the central Baltic Sea. *The ISME journal*, 4 (12), 1496-1508.

Lam, P., Jensen, M. M., Lavik, G., McGinnis, D. F., Müller, B., Schubert, C. J., Amann, R., Thamdrup, B., & Kuypers, M. M. (2007). Linking crenarchaeal and bacterial nitrification to anammox in the Black Sea. *Proceedings of the National Academy of Sciences*, 104 (17), 7104-7109.

Lam, P., Lavik, G., Jensen, M. M., van de Vossenberg, J., Schmid, M., Woebken, D., Gutl érez, D., Amann, R., Jetten, M., M., & Kuypers, M. M. (2009). Revising the

nitrogen cycle in the Peruvian oxygen minimum zone. *Proceedings of the National Academy of Sciences*, 106 (12), 4752-4757.

López - García, P., Moreira, D., López - López, A., & Rodríguez - Valera, F. (2001).

A novel haloarchaeal - related lineage is widely distributed in deep oceanic regions. *Environmental microbiology*, 3 (1), 72-78.

Lund, M. B., Smith, J. M., & Francis, C. A. (2012). Diversity, abundance and expression of nitrite reductase (nirK)-like genes in marine thaumarchaea. *The ISME journal*, 6 (10), 1966-1977.

Martens-Habbena, W., Berube, P. M., Urakawa, H., Jos é R., & Stahl, D. A. (2009).

Ammonia oxidation kinetics determine niche separation of nitrifying Archaea and Bacteria. *Nature*, 461 (7266), 976-979.

Massana, R., Murray, A. E., Preston, C. M., & DeLong, E. F. (1997). Vertical distribution and phylogenetic characterization of marine planktonic Archaea in the Santa Barbara Channel. *Applied and environmental microbiology*, 63 (1), 50-56.

Massana, R., DeLong, E. F., & Pedrós-Ali ó C. (2000). A few cosmopolitan phylotypes dominate planktonic archaeal assemblages in widely different oceanic provinces. *Applied and Environmental Microbiology*, 66 (5), 1777-1787.

Matsutani, N., Nakagawa, T., Nakamura, K., Takahashi, R., Yoshihara, K., & Tokuyama, T. (2011). Enrichment of a novel marine ammonia-oxidizing archaeon obtained from sand of an eelgrass zone. *Microbes and environments*, 26 (1), 23-29.

Merbt, S. N., Stahl, D. A., Casamayor, E. O., Mart í E., Nicol, G. W., & Prosser, J. I. (2012). Differential photoinhibition of bacterial and archaeal ammonia oxidation. *FEMS microbiology letters*, 327 (1), 41-46.

Mincer, T. J., Church, M. J., Taylor, L. T., Preston, C., Karl, D. M., & DeLong, E. F. (2007). Quantitative distribution of presumptive archaeal and bacterial nitrifiers in Monterey Bay and the North Pacific Subtropical Gyre. *Environmental microbiology*, 9 (5), 1162-1175.

Mosier, A. C., & Francis, C. A. (2008). Relative abundance and diversity of ammonia - oxidizing archaea and bacteria in the San Francisco Bay estuary. *Environmental microbiology*, 10 (11), 3002-3016.

Murray, A. E., Wu, K. Y., Moyer, C. L., Karl, D. M., & DeLong, E. F. (1999). Evidence for circumpolar distribution of planktonic Archaea in the Southern Ocean. *Aquatic Microbial Ecology*, 18 (3), 263.

Offre, P., Spang, A., & Schleper, C. (2013). Archaea in biogeochemical cycles. *Annual review of microbiology*, 67, 437-457.

Orphan, V. J., Hinrichs, K. U., Ussler, W. I. I. I., Paull, C. K., Taylor, L. T., Sylva, S. P., Hayes, J. M., & DeLong, E. F. (2001). Comparative analysis of methane-oxidizing archaea and sulfate-reducing bacteria in anoxic marine sediments. *Applied and Environmental Microbiology*, 67 (4), 1922-1934.

Ouverney, C. C., & Fuhrman, J. A. (2000). Marine planktonic archaea take up amino acids. *Applied and Environmental Microbiology*, 66 (11), 4829-4833.

Park, B. J., Park, S. J., Yoon, D. N., Schouten, S., Damsté J. S. S., & Rhee, S. K. (2010). Cultivation of autotrophic ammonia-oxidizing archaea from marine sediments in coculture with sulfur-oxidizing bacteria. *Applied and environmental microbiology*, 76 (22), 7575-7587.

Park, S. J., Kim, J. G., Jung, M. Y., Kim, S. J., Cha, I. T., Ghai, R., Martín-Cuarado, A., Rodríguez-Valera, F., & Rhee, S. K. (2012). Draft genome sequence of an ammonia-oxidizing archaeon, "Candidatus Nitrosopumilus sediminis" AR2, from Svalbard in the Arctic Circle. *Journal of bacteriology*, 194 (24), 6948-6949.

- Pedneault, E., Galand, P. E., Potvin, M., Tremblay, J. É., & Lovejoy, C. (2014). Archaeal amoA and ureC genes and their transcriptional activity in the Arctic Ocean. *Scientific reports*, 4.
- Pester, M., Schleper, C., & Wagner, M. (2011). The Thaumarchaeota: an emerging view of their phylogeny and ecophysiology. *Current opinion in microbiology*, 14 (3), 300-306.
- Pitcher, A., Villanueva, L., Hopmans, E. C., Schouten, S., Reichart, G. J., & Damsté J. S. S. (2011). Niche segregation of ammonia-oxidizing archaea and anammox bacteria in the Arabian Sea oxygen minimum zone. *The ISME journal*, 5 (12), 1896-1904.
- Pouliot, J., Galand, P. E., Lovejoy, C., & Vincent, W. F. (2009). Vertical structure of archaeal communities and the distribution of ammonia monooxygenase A gene variants in two meromictic High Arctic lakes. *Environmental Microbiology*, 11 (3), 687-699.
- Purkhold, U., Pommerening-Röser, A., Juretschko, S., Schmid, M. C., Koops, H. P., & Wagner, M. (2000). Phylogeny of all recognized species of ammonia oxidizers based on comparative 16S rRNA and amoA sequence analysis: implications for molecular diversity surveys. *Applied and Environmental Microbiology*, 66 (12), 5368-5382.

Purkhold, U., Wagner, M., Timmermann, G., Pommerening-Röser, A., & Koops, H. P. (2003). 16S rRNA and amoA-based phylogeny of 12 novel betaproteobacterial ammonia-oxidizing isolates: extension of the dataset and proposal of a new lineage within the nitrosomonads. *International Journal of Systematic and Evolutionary Microbiology*, 53 (5), 1485-1494.

Qin, W., Amin, S. A., Martens-Habbena, W., Walker, C. B., Urakawa, H., Devol, A. H., Ingalls, A. E., Moffett, J. W., Armbrust, E. V., & Stahl, D. A. (2014). Marine ammonia-oxidizing archaeal isolates display obligate mixotrophy and wide ecotypic variation. *Proceedings of the National Academy of Sciences*, 111 (34), 12504-12509.

Reigstad, L. J., Richter, A., Daims, H., Urich, T., Schwark, L., & Schleper, C. (2008). Nitrification in terrestrial hot springs of Iceland and Kamchatka. *FEMS Microbiology Ecology*, 64 (2), 167-174.

Rotthauwe, J. H., Witzel, K. P., & Liesack, W. (1997). The ammonia monooxygenase structural gene amoA as a functional marker: molecular fine-scale analysis of natural ammonia-oxidizing populations. *Applied and environmental microbiology*, 63 (12), 4704-4712.

Santoro, A. E., Francis, C. A., De Sieyes, N. R., & Boehm, A. B. (2008). Shifts in the relative abundance of ammonia - oxidizing bacteria and archaea across

physicochemical gradients in a subterranean estuary. *Environmental Microbiology*, 10 (4), 1068-1079.

Santoro, A. E., Casciotti, K. L., & Francis, C. A. (2010). Activity, abundance and diversity of nitrifying archaea and bacteria in the central California Current. *Environmental Microbiology*, 12 (7), 1989-2006.

Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., Lesniewski, R. A., Oakley, B. B., Parks, D. H., Robinson, C. J., Sahl, J. W., Stres, B., Thallinger, G. G., Van Horn, D. J., & Weber, C. F. (2009). Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Applied and environmental microbiology*, 75 (23), 7537-7541.

Shimada, K., Kamoshida, T., Itoh, M., Nishino, S., Carmack, E., McLaughlin, F., Zimmermann, S., & Proshutinsky, A. (2006). Pacific Ocean inflow: Influence on catastrophic reduction of sea ice cover in the Arctic Ocean. *Geophysical Research Letters*, 33 (8).

Sintes, E., Bergauer, K., De Corte, D., Yokokawa, T., & Herndl, G. J. (2013). Archaeal amoA gene diversity points to distinct biogeography of ammonia - oxidizing Crenarchaeota in the ocean. *Environmental microbiology*, 15 (5), 1647-1658.



Smith, J. M., Casciotti, K. L., Chavez, F. P., & Francis, C. A. (2014). Differential contributions of archaeal ammonia oxidizer ecotypes to nitrification in coastal surface waters. *The ISME journal*, 8 (8), 1704-1714.

Stein, J. L., & Simon, M. I. (1996). Archaeal ubiquity. *Proceedings of the National Academy of Sciences of the United States of America*, 93 (13), 6228.

Spang, A., Hatzenpichler, R., Brochier-Armanet, C., Rattei, T., Tischler, P., Spieck, E., Streit, W., Stahl, D. A., Wagner, M., & Schleper, C. (2010). Distinct gene set in two different lineages of ammonia-oxidizing archaea supports the phylum Thaumarchaeota. *Trends in microbiology*, 18 (8), 331-340.

Suzuki, M. T., Taylor, L. T., & DeLong, E. F. (2000). Quantitative analysis of small-subunit rRNA genes in mixed microbial populations via 5'-nuclease assays. *Applied and Environmental Microbiology*, 66 (11), 4605-4614.

Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., & Kumar, S. (2011). MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular biology and evolution*, 28 (10), 2731-2739.

Taylor, A. E., & Bottomley, P. J. (2006). Nitrite production by *Nitrosomonas europaea* and *Nitrosospira* sp. AV in soils at different solution concentrations of ammonium. *Soil Biology and Biochemistry*, 38 (4), 828-836.

Tolar, B. B., King, G. M., & Hollibaugh, J. T. (2013). An analysis of Thaumarchaeota populations from the northern Gulf of Mexico. *Frontiers in microbiology*, 4, DOI: 10.3389/fmicb.2013.00072

Trias, R., Garc ía-Lled ó, A., S áncchez, N., L ópez-Jurado, J. L., Hallin, S., & Ba ñeras, L. (2012). Abundance and composition of epiphytic bacterial and archaeal ammonia oxidizers of marine red and brown macroalgae. *Applied and environmental microbiology*, 78 (2), 318-325.

Urakawa, H., Tajima, Y., Numata, Y., & Tsuneda, S. (2008). Low temperature decreases the phylogenetic diversity of ammonia-oxidizing archaea and bacteria in aquarium biofiltration systems. *Applied and environmental microbiology*, 74 (3), 894-900.

Venter, J. C., Remington, K., Heidelberg, J. F., Halpern, A. L., Rusch, D., Eisen, J. A., Wu, D., Paulsen, I., Nelson, K. E., Nelson, W., Fouts, D. E., Levy, S., Knap, A. H., Lomas, M. W., Nealson, K., White, O., Peterson, J., Hoffman, J., Parsons, R., Baden-

Tillson, H., Pfannkoch, C., Rogers, Y., & Smith, H. O. (2004). Environmental genome shotgun sequencing of the Sargasso Sea. *science*, 304 (5667), 66-74.

Verhamme, D. T., Prosser, J. I., & Nicol, G. W. (2011). Ammonia concentration determines differential growth of ammonia-oxidising archaea and bacteria in soil microcosms. *The ISME journal*, 5 (6), 1067-1071.

Walker, C. B., De La Torre, J. R., Klotz, M. G., Urakawa, H., Pinel, N., Arp, D. J., Brochier-Armanet, C., Chain, P. S., Chan, P. P., Gollabqir, A., Hemp, J., Hügler, M., Karr, E. A., Könneke, M., Shin, M., Lawton, T. J., Lowe, T., Martens-Habbena, W., Sayavedra-Soto, L. A., Lang, D., Sievert, S. M., Rosenzweig, A. C., Manning, G., & Stahl, D. A. (2010). Nitrosopumilus maritimus genome reveals unique mechanisms for nitrification and autotrophy in globally distributed marine crenarchaea. *Proceedings of the National Academy of Sciences*, 107 (19), 8818-8823.

Wang, S., Xiao, X., Jiang, L., Peng, X., Zhou, H., Meng, J., & Wang, F. (2009). Diversity and abundance of ammonia-oxidizing archaea in hydrothermal vent chimneys of the Juan de Fuca Ridge. *Applied and environmental microbiology*, 75 (12), 4216-4220.

Ward, B. B., Martino, D. P., Diaz, M. C., & Joye, S. B. (2000). Analysis of ammonia-oxidizing bacteria from hypersaline Mono Lake, California, on the basis of 16S rRNA sequences. *Applied and environmental microbiology*, 66 (7), 2873-2881.

Ward, B. B., & O'Mullan, G. D. (2002). Worldwide distribution of *Nitrosococcus oceanus*, a marine ammonia-oxidizing  $\gamma$ -proteobacterium, detected by PCR and sequencing of 16S rRNA and amoA genes. *Applied and environmental microbiology*, 68 (8), 4153-4157.

Walsh, J. J., Dieterle, D. A., Maslowski, W., & Whitledge, T. E. (2004). Decadal shifts in biophysical forcing of Arctic marine food webs: numerical consequences. *Journal of Geophysical Research: Oceans* (1978–2012), 109(C5).

Williams, W. J., Shroyer, E., Kinney, J. C., Itoh, M., & Maslowski, W. (2014). Shelf-break exchange in the Bering, Chukchi and Beaufort Seas. In *The Pacific Arctic Region* (pp. 133-165). Springer Netherlands.

Woese, C. R., & Fox, G. E. (1977). Phylogenetic structure of the prokaryotic domain: the primary kingdoms. *Proceedings of the National Academy of Sciences*, 74 (11), 5088-5090.

Woese, C. R., Kandler, O., & Wheelis, M. L. (1990). Towards a natural system of organisms: proposal for the domains Archaea, Bacteria, and Eucarya. *Proceedings of the National Academy of Sciences*, 87 (12), 4576-4579.

Woodgate, R. A., Aagaard, K., & Weingartner, T. J. (2005). Monthly temperature, salinity, and transport variability of the Bering Strait through flow. *Geophysical Research Letters*, 32 (4).

Wuchter, C., Abbas, B., Coolen, M. J., Herfort, L., van Bleijswijk, J., Timmers, P., Strous, M., Teira, E., Herndl, G. J., Middelburg, J. J., Schouten, S., & Damsté J. S. S. (2006). Archaeal nitrification in the ocean. *Proceedings of the National Academy of Sciences*, 103 (33), 12317-12322.

Yakimov, M. M., La Cono, V., Smedile, F., DeLuca, T. H., Juárez, S., Ciordia, S., Fernández, M., Albar, J. P., Ferrer, M., Golyshin, P. N., & Giuliano, L. (2011). Contribution of crenarchaeal autotrophic ammonia oxidizers to the dark primary production in Tyrrhenian deep waters (Central Mediterranean Sea). *The ISME journal*, 5 (6), 945-961.